

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 May 2002 (10.05.2002)

PCT

(10) International Publication Number
WO 02/36131 A1(51) International Patent Classification⁷: A61K 31/70,
A01N 43/04, C12N 15/63, 15/86

(21) International Application Number: PCT/US01/42885

(22) International Filing Date:
1 November 2001 (01.11.2001)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/244,598 1 November 2000 (01.11.2000) US(71) Applicants: LUDWIG INSTITUTE FOR CANCER
RESEARCH [US/US]; 605 Third Avenue, New York, NY
10158 (US). AVENTIS PHARMA S.A. [FR/FR]; 20 Avenue
Raymond Aron, F-92160 Antony (FR). LICENTIA
LTD. [FI/FI]; Erottajankatu 19 B 5, FIN-00130 Helsinki
(FI).(72) Inventors: ENHOLM, Berndt; Haartman Institute,
University of Helsinki, P.O. Box 21, FIN-00014 (FI).
BRANELLEC, Didier; 13 Quai Jules Guesde, BP 14,
F-944403 Vitry-sur-Seine (FR). YLAIE-HERTTUALA,Seppo; A. I. Virtanen Institute, University of Kuopio, P.O.
Box 1627 (US). PETTERSSON, Ralf; Ludwig Institute
for Cancer Research, Karolinska Institute, Box 240, S-171
77 Stockholm (SE). BERGSTEN, Erika; Ludwig Institute
for Cancer Research, Karolinska Institute, Box 240, S-171
77 Stockholm (SE). ERIKSSON, Ulf; Ludwig Institute
for Cancer Research, Karolinska Institute, Box 240, S-171
77 Stockholm (SE). ALITALO, Karl; Haartman Institute,
University of Helsinki, P.O. Box 21, SF-00014 Helsinki
(FI). LE ROUX, Aude; 13, allée Chenard et Walcker,
F-94320 Thiais (FR).(74) Agent: EVANS, Joseph, D.; Crowell & Moring LLP, Intel-
lectual Property Group, P.O. Box 14300, Washington, DC
20044-4300 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian

[Continued on next page]

(54) Title: IN VIVO STIMULATION OF ANGIOGENIC ACTIVITY

(57) Abstract: Angiogenesis is stimulated by introducing two replication-deficient adenovirus vectors to the endothelial cells or proximate to the endothelial cells of an organism. The first vector encodes VEGF-B₁₆₇ or a fragment or conservative substitution thereof, the second vector encodes VEGF-A or VEGF-C, or fragments or conservative substitutions thereof.



patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

IN VIVO STIMULATION OF ANGIOGENIC ACTIVITY

[0001] This application claims the priority of U.S. Provisional Patent Application No. 60/244,598, filed on November 1, 2000, the disclosure of which is expressly incorporated by reference herein.

BACKGROUND OF THE INVENTION

[0002] The present invention relates generally to gene therapy, more specifically, to virus-mediated and other forms of gene therapy. More particularly, the invention relates to adenovirus-mediated delivery of genes useful in the promotion of angiogenesis.

[0003] Angiogenesis is the formation of new capillary blood vessels by a process of sprouting from pre-existing vessels, and occurs during development as well as in a number of physiological and pathological processes. Angiogenesis is a physiologically complex process involving proliferation of endothelial cells, degradation of extracellular matrix, branching of vessels and subsequent cell adhesion events. In the adult, angiogenesis is tightly controlled and limited under normal circumstances to the female reproductive system. Angiogenesis can, however, be switched on in response to tissue damage. Solid tumors are also able to induce angiogenesis in surrounding tissue, thus sustaining tumor growth and facilitating the formation of metastases (Folkman, J., Nature Med. 1:27-31, (1995)). The molecular mechanisms underlying the complex angiogenic processes are far from being understood. A similar although far less well studied process also occurs in the lymphatic system, and is sometimes referred to as lymphangiogenesis.

[0004] Angiogenesis begins with localized breakdown of the basement membrane of the parent vessel, which is followed by the migration and outgrowth of endothelial cells into the surrounding extracellular matrix, resulting in the formation of a capillary sprout. A lumen is subsequently formed,

and constitutes an essential element in functional sprout formation. Sprout maturation is completed after reconstitution of the basement membrane.

[0005] Alterations in at least three endothelial cell functions occur during this series of events: 1) modulation of interactions with the extracellular matrix, which requires alterations in cell-matrix contacts and the production of matrix-degrading proteolytic enzymes (plasminogen activators (PAs) and matrix metalloproteinases); 2) an initial increase and subsequent decrease in locomotion (migration), which allows the cells to translocate towards the angiogenic stimulus and to stop once they reach their destination; 3) an increase in proliferation, which provides new cells for the growing and elongating vessel, and a subsequent return to the quiescent state once the vessel is formed.

[0006] Together, these cellular functions contribute to the process of capillary morphogenesis, i.e. the formation of three-dimensional patent or open tube-like structures. Many newly formed capillaries subsequently undergo a process of vessel wall maturation (i.e. formation of a smooth muscle cell-rich media and an adventitia, while others undergo regression (i.e. in the absence of blood flow) [see Pepper *et al.*, *Enzyme Protein*, 49:138-162 (1996); Risau, *Nature* 386:671-674 (1997)].

[0007] With the exception of angiogenesis which occurs in response to tissue injury or in female reproductive organs, endothelial cell turnover in the healthy adult organism is very low. The maintenance of endothelial quiescence is thought to be due to the presence of endogenous negative regulators, since positive regulators are frequently detected in adult tissues in which there is apparently no angiogenesis. The converse is also true, namely that positive and negative regulators often co-exist in tissues in which endothelial cell turnover is increased. This has lead to the notion of the "angiogenic switch", in which endothelial activation status is determined by a balance between positive and negative regulators.

[0008] In activated (angiogenic) endothelium, positive regulators predominate, whereas endothelial quiescence is achieved and maintained by the dominance of negative regulators [Hanahan *et al.*, *Cell*, 86:353-364 (1996)]. Used initially in the context of tumor progression to describe the passage from the

prevascular to the vascular phase, the notion of the "switch" can also be applied in the context of developmental, physiological as well as pathological angiogenesis. Although it still remains to be definitively demonstrated *in vivo*, the current working hypothesis is that the "switch" involves the induction of a positive regulator and/or the loss of a negative regulator.

[0009] Because of the crucial role of angiogenesis in so many physiological and pathological processes, factors involved in the control of angiogenesis have been intensively investigated. A number of growth factors have been shown to be involved in the regulation of angiogenesis; these include fibroblast growth factors (FGFs), platelet-derived growth factors (PDGFs), transforming growth factor alpha (TGF-alpha), and hepatocyte growth factor (HGF). See for example Folkman *et al.*, *J. Biol. Chem.*, 267:10931-10934 (1992) for a review.

[0010] It has been suggested that a particular family of endothelial cell-specific growth factors, the vascular endothelial growth factors (VEGFs), and their corresponding receptors is primarily responsible for stimulation of endothelial cell growth and differentiation, and for certain functions of the differentiated cells. These factors are members of the PDGF/VEGF family, and appear to act primarily via endothelial receptor tyrosine kinases (RTKs). The PDGF/VEGF family of growth factors belongs to the cystine-knot superfamily of growth factors, which also includes the neurotrophins and transforming growth factor- β .

[0011] Eight different proteins have been identified in the PDGF/VEGF family, namely two PDGFs (A and B), VEGF-A and five members that are closely related to VEGF-A. The five members closely related to VEGF-A are: VEGF-B, described in International Patent Application PCT/US96/02957 (WO 96/26736) and in U.S. Patents 5,840,693 and 5,607,918 by Ludwig Institute for Cancer Research and The University of Helsinki; VEGF-C or VEGF2, described in Joukov *et al.*, *EMBO J.* 15:290-298 (1996), Lee *et al.*, *Proc. Natl. Acad. Sci. USA* 93:1988-1992 (1996), and U.S. Patents 5,932,540 and 5,935,540 by Human Genome Sciences, Inc; VEGF-D, described in International Patent Application No. PCT/US97/14696 (WO 98/07832), and Achen *et al.*, *Proc. Natl. Acad. Sci. USA* 95:548-553 (1998); the placenta growth factor (PlGF), described in Maglione

et al., *Proc. Natl. Acad. Sci. USA* 88:9267-9271 (1991); and VEGF3, described in International Patent Application No. PCT/US95/07283 (WO 96/39421) by Human Genome Sciences, Inc.

[0012] Each VEGF family member has between 30% and 45% amino acid sequence identity with VEGF-A in their VEGF homology domain (VHD). This VHD contains the eight conserved cysteine residues which form the cystine-knot motif. In their active, physiological state, the proteins are dimers. Functional characteristics of the VEGF family include varying degrees of mitogenicity for endothelial cells and related cell types, induction of vascular permeability and angiogenic and lymphangiogenic properties.

[0013] Vascular endothelial growth factor (VEGF-A) is a homodimeric glycoprotein that has been isolated from several sources. VEGF-A shows highly specific mitogenic activity for endothelial cells. VEGF-A has important regulatory functions in the formation of new blood vessels during embryonic vasculogenesis and in angiogenesis during adult life (Carmeliet *et al.*, *Nature*, 380: 435-439, (1996); Ferrara *et al.*, *Nature*, 380: 439-442, (1996); reviewed in Ferrara and Davis-Smyth, *Endocrine Rev.*, 18: 4-25, (1997)). The significance of the role played by VEGF-A has been demonstrated in studies showing that inactivation of a single VEGF-A allele results in embryonic lethality due to failed development of the vasculature (Carmeliet *et al.*, *Nature*, 380: 435-439, (1996); Ferrara *et al.*, *Nature*, 380: 439-442, (1996)). The isolation and properties of VEGF-A have been reviewed; see Ferrara *et al.*, *J. Cellular Biochem.*, 47: 211-218, (1991) and Connolly, *J. Cellular Biochem.*, 47: 219-223, (1991).

[0014] In addition VEGF-A has strong chemoattractant activity towards monocytes, can induce the plasminogen activator and the plasminogen activator inhibitor in endothelial cells, and can also induce microvascular permeability. Because of the latter activity, it is sometimes referred to as vascular permeability factor (VPF). VEGF-A is also chemotactic for certain hematopoietic cells. Recent literature indicates that VEGF-A blocks maturation of dendritic cells and thereby reduces the effectiveness of the immune response to tumors (many tumors secrete VEGF-A) (Gabrilovich *et al.*, *Blood* 92: 4150-4166, (1998); Gabrilovich *et al.*, *Clinical Cancer Research* 5: 2963-2970, (1999)).

[0015] Vascular endothelial growth factor B (VEGF-B) has similar angiogenic and other properties to those of VEGF-A, but is distributed and expressed in tissues differently from VEGF-A. In particular, VEGF-B is very strongly expressed in heart, and only weakly in lung, whereas the reverse is the case for VEGF-A (Olofsson, B. *et al.*, *Proc. Natl. Acad. Sci. USA* 93:2576-2581 (1996)). RT-PCR assays have demonstrated the presence of VEGF-B mRNA in melanoma, normal skin, and muscle. This suggests that VEGF-A and VEGF-B, despite the fact that they are co-expressed in many tissues, may have functional differences.

[0016] A comparison of the PDGF/VEGF family of growth factors reveals that the 167 amino acid isoform of VEGF-B is the only family member that is completely devoid of any glycosylation. Gene targeting studies have shown that VEGF-B deficiency results in mild cardiac phenotype, and impaired coronary vasculature (Bellomo *et al.*, *Circ. Res.* 86:E29-35 (2000)). VEGF-B knock out mice were demonstrated to have impaired coronary vessel structure, smaller hearts and impaired recovery after cardiac ischemia (Bellomo, D. *et al.*, *Circulation Research (Online)*, 86:E29-35 (2000)).

[0017] Human VEGF-B was isolated using a yeast co-hybrid interaction trap screening technique by screening for cellular proteins which might interact with cellular retinoic acid-binding protein type I (CRABP-I). The isolation and characteristics including nucleotide and amino acid sequences for both the human and mouse VEGF-B are described in detail in PCT/US96/02957, in U.S. Patents 5,840,693 and 5,607,918 by Ludwig Institute for Cancer Research and The University of Helsinki and in Olofsson *et al.*, *Proc. Natl. Acad. Sci. USA* 93:2576-2581 (1996). The nucleotide sequence for human VEGF-B is also found at GenBank Accession No. U48801. The entire disclosures of the International Patent Application PCT/US97/14696 (WO 98/07832), U.S. Patents 5,840,693 and 5,607,918 are incorporated herein by reference.

[0018] The mouse and human genes for VEGF-B are almost identical, and both span about 4 kb of DNA. The genes are composed of seven exons and their exon-intron organization resembles that of the VEGF and PlGF genes (Grimmond *et al.*, *Genome Res.* 6:124-131 (1996); Olofsson *et al.*, *J. Biol. Chem.*

271:19310-19317 (1996); Townson *et al.*, *Biochem. Biophys. Res. Commun.* 220:922-928 (1996)).

[0019] VEGF-C was isolated from conditioned media of the PC-3 prostate adenocarcinoma cell line (CRL1435) by screening for ability of the medium to induce tyrosine phosphorylation of the endothelial cell-specific receptor tyrosine kinase VEGFR-3 (Flt4), using cells transfected to express VEGFR-3. VEGF-C was purified using affinity chromatography with recombinant VEGFR-3, and was cloned from a PC-3 cDNA library. Its isolation and characteristics are described in detail in Joukov *et al.*, *EMBO J.*, 15: 290-298, (1996).

[0020] VEGF-D was isolated from a human breast cDNA library, commercially available from Clontech, by screening with an expressed sequence tag obtained from a human cDNA library designated "Soares Breast 3NbHBst" as a hybridization probe (Achen *et al.*, *Proc. Natl. Acad. Sci. USA*, 95: 548-553, (1998)). Its isolation and characteristics are described in detail in International Patent Application No. PCT/US97/14696 (WO98/07832). In PCT/US97/14696, the isolation of a biologically active fragment of VEGF-D is also described. This fragment consists of VEGF-D amino acid residues 93 to 201.

[0021] The VEGF-D gene is broadly expressed in the adult human, but is certainly not ubiquitously expressed. VEGF-D is strongly expressed in heart, lung and skeletal muscle. Intermediate levels of VEGF-D are expressed in spleen, ovary, small intestine and colon, and a lower expression occurs in kidney, pancreas, thymus, prostate and testis. No VEGF-D mRNA was detected in RNA from brain, placenta, liver or peripheral blood leukocytes.

[0022] PlGF was isolated from a term placenta cDNA library. Its isolation and characteristics are described in detail in Maglione *et al.*, *Proc. Natl. Acad. Sci. USA*, 88: 9267-9271, (1991). Presently its biological function is not well understood.

[0023] VEGF3 was isolated from a cDNA library derived from colon tissue. VEGF3 is stated to have about 36% identity and 66% similarity to VEGF. The method of isolation of the gene encoding VEGF3 is unclear and no characterization of the biological activity is disclosed in International Patent Application No. PCT/US95/07283 (WO 96/39421).

[0024] Similarity between two proteins is determined by comparing the amino acid sequence and conserved amino acid substitutions of one of the proteins to the sequence of the second protein, whereas identity is determined without including the conserved amino acid substitutions.

[0025] As noted above, the PDGF/VEGF family members act primarily by binding to receptor tyrosine kinases. In general, receptor tyrosine kinases are glycoproteins, which consist of an extracellular domain capable of binding a specific growth factor(s), a transmembrane domain, which is usually an alpha-helical portion of the protein, a juxtamembrane domain, which is where the receptor may be regulated by, *e.g.*, protein phosphorylation, a tyrosine kinase domain, which is the enzymatic component of the receptor and a carboxy-terminal tail, which in many receptors is involved in recognition and binding of the substrates for the tyrosine kinase.

[0026] Five endothelial cell-specific receptor tyrosine kinases have been identified, belonging to two distinct subclasses: three vascular endothelial cell growth factor receptors, VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt4), and the two receptors of the Tie family, Tie and Tie-2 (Tek). These receptors differ in their specificity and affinity. All of these have the intrinsic tyrosine kinase activity which is necessary for signal transduction.

[0027] The only receptor tyrosine kinases known to bind VEGFs are VEGFR-1, VEGFR-2 and VEGFR-3. VEGFR-1 and VEGFR-2 bind VEGF with high affinity, and VEGFR-1 also binds PlGF. VEGF-B binds to VEGFR-1 with high affinity, but not to VEGFR-2 or -3 (Olofsson *et al.*, *Proc. Natl. Acad. Sci. USA*, 95:11709-11714 (1998)). VEGF-C has been shown to be the ligand for VEGFR-3, and it also activates VEGFR-2 (Joukov *et al.*, *EMBO J.* 15:290-298 (1996)). VEGF-D binds to both VEGFR-2 and VEGFR-3 (Achen *et al.*, *Proc. Natl. Acad. Sci. USA* 95:548-553 (1998)). A ligand for Tek/Tie-2 has been described in International Patent Application No. PCT/US95/12935 (WO 96/11269) by Regeneron Pharmaceuticals, Inc. The ligand for Tie has not yet been identified.

[0028] A novel 130-135 kDa VEGF isoform specific receptor has been purified and cloned (Soker *et al.*, *Cell* 92:735-745 (1998)). The VEGF receptor was found to specifically bind the VEGF₁₆₅ isoform via the exon 7 encoded

sequence, which shows weak affinity for heparin (Soker *et al.*, *Cell* 92:735-745 (1998)). Surprisingly, the receptor was shown to be identical to human neuropilin-1 (NP-1), a receptor involved in early stage neuromorphogenesis. VEGF-B₁₆₇ also binds to NP-1 (Makinen, T. *et al.*, *J Biol Chem.*, 274:21217-21222(1999)). In addition, PlGF-2 also appears to interact with NP-1 (Migdal *et al.*, *J. Biol. Chem.* 273:22272-22278 (1998)).

[0029] VEGFR-1, VEGFR-2 and VEGFR-3 are expressed differently by endothelial cells. Generally, both VEGFR-1 and VEGFR-2 are expressed in blood vessel endothelia (Oelrichs *et al.*, *Oncogene* 8:11-18 (1992); Kaipainen *et al.*, *J. Exp. Med.* 178:2077-2088 (1993); Dumont *et al.*, *Dev. Dyn.* 203:80-92 (1995); Fong *et al.*, *Dev. Dyn.* 207:1-10 (1996)) and VEGFR-3 is mostly expressed in the lymphatic endothelium of adult tissues (Kaipainen *et al.*, *Proc. Natl. Acad. Sci. USA* 9:3566-3570 (1995)). VEGFR-3 is also expressed in the blood vasculature surrounding tumors.

[0030] VEGFRs are expressed in many adult tissues, despite the apparent lack of constitutive angiogenesis. VEGFRs are however clearly upregulated in endothelial cells during development and in certain angiogenesis-associated/dependent pathological situations including tumor growth [see Dvorak *et al.*, *Amer. J. Pathol.*, 146:1029-1039 (1995); Ferrara *et al.*, *Endocrine Rev.*, 18:4-25 (1997)]. The phenotypes of VEGFR-1-deficient mice and VEGFR-2-deficient mice reveal an essential role for these receptors in blood vessel formation during development.

[0031] VEGFR-1-deficient mice die *in utero* at mid-gestation due to inefficient assembly of endothelial cells into blood vessels, resulting in the formation of abnormal vascular channels [Fong *et al.*, *Nature*, 376:66-70 (1995)]. VEGFR-2-deficient mice die *in utero* between 8.5 and 9.5 days post-coitum, and in contrast to VEGFR-1, this appears to be due to abortive development of endothelial cell precursors [Shalaby *et al.*, *Nature* 376:62-66 (1995)]. The importance of VEGFR-2 in tumor angiogenesis has also been clearly demonstrated by using a dominant-negative approach [Millauer *et al.*, *Nature*, 367:576-579 (1994); Millauer *et al.*, *Cancer Res.* 56:1615-1620 (1996)]. The phenotype of VEGFR-3-deficient mice has been reported in Dumont, *et al.*,

Cardiovascular Failure in Mouse Embryos Deficient in VEGF Receptor-3, *Science*, 282:946-949 (1998). VEGFR-3 deficient mice die *in utero* between 12 and 14 days of gestation due to defective blood vessel development.

[0032] Despite intensive activity in the art, there has remained a need for ways to stimulate angiogenesis *in vivo*.

SUMMARY OF THE INVENTION

[0033] The present invention is directed to a gene therapy approach useful in stimulation of angiogenesis. One objective of the present invention is to provide a method for stimulating angiogenesis in which VEGF-B₁₆₇ proteins or peptides in addition to other VEGF proteins and peptides as are produced to a therapeutically significant degree for sustained periods to stimulate angiogenesis with at least a first vector construct having a polynucleotide sequence for VEGF-B₁₆₇ and a second vector construct containing a polynucleotide sequence for another VEGF, preferably a replication-deficient adenovirus construct.

[0034] According to one embodiment of the present invention, a method of stimulating angiogenesis comprises introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism and introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-A or a fragment or conservative substitution thereof to the organism. In this embodiment the replication-deficient adenovirus vectors are delivered to at least one cell of the organism, the target cell being either an endothelial cell or a cell in proximity to an endothelial cell.

[0035] According to a further embodiment, the target cells are vascular cells. According to a further embodiment, the target cells are either microvascular endothelial cells or aortic endothelial cells.

[0036] According to a further embodiment, the organism may be mammalian, murine, or human.

[0037] According to a further embodiment, 10^7 to 10^{13} of vector particles of each adenovirus vector are introduced to the organism.

[0038] According to a further embodiment, expression of the polynucleotide sequence is driven by a CMV promoter which is contained in the vector.

[0039] According to a further embodiment of the present invention, a method of stimulating angiogenesis comprises introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism and introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-C or a fragment or conservative substitution thereof to the organism. In this embodiment the replication-deficient adenovirus vectors are delivered to at least one cell of the organism, the target cell being either an endothelial cell or a cell in proximity to an endothelial cell.

[0040] According to a further embodiment of the present invention, a method of stimulating angiogenesis comprises introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism and introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-D or a fragment or conservative substitution thereof to the organism. In this embodiment the replication-deficient adenovirus vectors are delivered to at least one cell of the organism, the target cell being either an endothelial cell or a cell in proximity to an endothelial cell.

[0041] The reasons for using a replication-deficient adenovirus construct are that it allows efficient *in vivo* expression of the transgene. The virus is produced to high titers and many cell types can be infected with adenovirus. It also allows for the simultaneous expression of more than one transgene by coinfection with the recombinant adenoviruses (e.g. VEGF-B + VEGF-A). Expression is also very fast, usually within 24-48 hours.

[0042] The method involves stimulating angiogenesis by delivering VEGF-B₁₆₇ and another VEGF to target cells of an organism in need of such stimulation. The target cells include, but are not limited to, endothelial cells or

cells in the proximity of endothelial cells, vascular endothelial cells, microvascular endothelial cells and aortic endothelial cells. These cells can be from a mammalian organism, such as murine or human cells. Preferably the expression of the polynucleotide sequences is driven by a cytomegalovirus (CMV) promoter. Preferably about 10^7 to about 10^{13} of vector particles of each adenovirus vector are delivered *in vivo*.

[0043] Because the nude mouse has been proven to be a relevant animal model for angiogenesis, a similar effect can be expected in humans.

[0044] These recombinant adenoviruses can be administered as a pharmaceutical composition in combination with any standard physiologically and/or pharmaceutically acceptable carriers known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue or organism. The characteristics of the carrier will depend on the route of administration. Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art.

[0045] It is within the scope of the present invention to employ functional fragments of the desired sequences. Techniques known in the art may be employed to conserve the regions of the sequence necessary for proper function, while performing appropriate substitutions and/or deletions to the remaining portions of the sequence so that the resultant product maintains the function of the full sequence.

[0046] As used herein, the term "conservative substitution" denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative substitutions include the substitution of one hydrophobic residue such as isoleucine, valine, leucine, alanine, cysteine, glycine, phenylalanine, proline, tryptophan, tyrosine, norleucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic acid for aspartic acid, or glutamine

for asparagine, and the like. Neutral hydrophilic amino acids which can be substituted for one another include asparagine, glutamine, serine and threonine. The term "conservative substitution" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid.

[0047] As such, it should be understood that in the context of the present invention, a conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in the following Table A from WO 97/09433.

Table A
Conservative Substitutions I

SIDE CHAIN	
CHARACTERISTIC	AMINO ACID
Aliphatic	
Non-polar	G A P I L V
Polar - uncharged	C S T M N Q
Polar - charged	D E K R
Aromatic	H F W Y
Other	N Q D E

[0048] Alternatively, conservative amino acids can be grouped as described in Lehninger, [Biochemistry, Second Edition; Worth Publishers, Inc. NY:NY (1975), pp.71-77] as set out in the following Table B.

Table B

Conservative Substitutions II

SIDE CHAIN
CHARACTERISTIC AMINO ACID

Non-polar (hydrophobic)	
A. Aliphatic:	A L I V P
B. Aromatic:	F W
C. Sulfur-containing:	M
D. Borderline:	G
Uncharged-polar	
A. Hydroxyl:	S T Y
B. Amides:	N Q
C. Sulfhydryl:	C
D. Borderline:	G
Positively Charged (Basic):	K R H
Negatively Charged (Acidic):	D E

[0049] Exemplary conservative substitutions are set out in the following Table C.

Table C
Conservative Substitutions III

<u>Original Residue</u>	<u>Exemplary Substitution</u>
Ala (A)	Val, Leu, Ile
Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg
Ile (I)	Leu, Val, Met, Ala, Phe,
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	Gly
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr, Phe
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

[0050] If desired, the peptides of the invention can be modified, for instance, by glycosylation, amidation, carboxylation, or phosphorylation, or by the creation of acid addition salts, amides, esters, in particular C-terminal esters, and N-acyl derivatives of the peptides of the invention. The peptides also can be modified to create peptide derivatives by forming covalent or noncovalent

complexes with other moieties. Covalently-bound complexes can be prepared by linking the chemical moieties to functional groups on the side chains of amino acids comprising the peptides, or at the N- or C-terminus.

[0051] It is anticipated that the aforementioned peptides can be conjugated to a reporter group, including, but not limited to a radiolabel, a fluorescent label, an enzyme (e.g., that catalyzes a colorimetric or fluorometric reaction), a substrate, a solid matrix, or a carrier (e.g., biotin or avidin).

BRIEF DESCRIPTION OF THE DRAWINGS

[0052] The invention will be described in further detail hereinafter with reference to the accompanying drawings in which:

[0053] Fig. 1 represents a blot of an immunoprecipitation of 293 EBNA cells infected with AdVEGF-B₁₆₇ using polyclonal antibodies against VEGF-B;

[0054] Figs. 2A-2F show results of stained sections of the skin of mouse ears injected subcutaneously with the recombinant adenoviruses:

[0055] Figs. 2B and 2E-AdVEGF,

[0056] Figs 2A and 2D-AdLacZ,

[0057] Figs 2C and 2F-AdVEGF-A + AdVEGF-B; and

[0058] Fig. 3 provides quantitative results of blood vessel formation.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Example 1: Production of Recombinant Adenovirus with VEGF-B₁₆₇ or VEGF-A

[0059] To make AdVEGF-B₁₆₇ vector, the full-length VEGF-B₁₆₇ cDNA was amplified from a plasmid by PCR (forward primer: 5' CGA TCT GGC CAT ACA CTT 3' (SEQ ID NO:1); reverse primer 5' CTA TGG ATC CTC ACC TTC GCA GCT T 3' (SEQ ID NO:2), contains a BamHI site) and then cloned into the TA-plasmid pCR2.1 vector (Invitrogen).

[0060] Following amplification in *E. coli* and plasmid purification, the insert was excised with BamHI and cloned into the BamHI site of the plasmid pQBI-AdCMV5-IRES-GFP (Quantum Biotechnologies, Quebec, Canada; transfer

plasmid). In this plasmid the cDNA expression is driven from the CMV promoter. The plasmid also contains an IRES-element allowing the simultaneous expression of green fluorescent protein (GFP). Expression of VEGF-B was verified by transfecting the plasmid into COS-1 cells.

[0061] The above transfer plasmid was linearized with FseI and then cotransfected with pJM17, a circular adenovirus plasmid (Microbix Biosystems Inc, Ontario, Canada), into 293 cells. Transfection was with Lipofectamine Plus. Fourteen days later, cells were harvested, broken by freeze-thawing, and the supernatant from a low-speed centrifugation was used to infect new 293 cells. The amplification step was repeated once. Expression of the insert was verified using GFP as an indicator. After this, the recombinant virus was plaque-purified in 293 cells once using standard protocols. A GFP-positive plaque was amplified and subsequently used to prepare a stock that was purified by CsCl centrifugation.

[0062] To make the AdVEGF-A vector, full-length VEGF-A was cloned into a first-generation adenovirus vector described in Hiltunen *et al.*, *Circulation*, 102:2262-2268 (2000). This reference is expressly incorporated by reference. Alternatively, active fragments of VEGF-A or either the entire or fragmented sequence containing conservative substitutions could be employed. The preparation of other vectors and the use of different vector preparation techniques known in the art are within the scope of the present invention. Further, the techniques employed to create the VEGF-B₁₆₇ or VEGF-A adenovirus vectors could be employed to create VEGF-C or VEGF-D adenovirus vectors.

[0063] 293 EBNA cells were incubated for one hour in serum free media containing 2×10^7 pfu of AdVEGF-B₁₆₇. Additionally, cells were transfected with pREP7 (Invitrogen) plasmid expression vectors encoding VEGF-B or pREP7 vector control using the calcium phosphate method. Cells were metabolically labeled the next day with S³⁵-methionine and cysteine (Promix, Amersham) for 6 hours. The media were collected and labeled VEGF-B proteins were precipitated using a polyclonal antibody raised against VEGF-B [see Olofsson, B. *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:2576-2581 (1996)]. The precipitated proteins bound to

protein A sepharose were washed three times in phosphate-buffered saline, dissolved in Laemmli sample buffer and analyzed on a 12.5% or 15% SDS-PAGE gel. Gels were then dried and analysed by a phosphoimager and autoradiography. As seen in Figure 1, cells infected with the AdVEGF-B₁₆₇ or transfected with the pREC7 plasmids encoding VEGF-B₁₆₇ produced major polypeptides of about 21 kD.

Example 2: Evaluation of Angiogenic Response to Recombinant Adenoviruses

[0064] About 2×10^8 pfu of each of the VEGF-B₁₆₇ and VEGF-A recombinant adenoviruses were injected subcutaneously into five ears of three NMRI nu/nu mice (Harlan, the Netherlands). The mice were sacrificed at various time points after infection and the skin from the site of injection was fixed in 4% paraformaldehyde and embedded in paraffin. Seven micrometer sections were stained with monoclonal antibodies against VEGFR-2 [Kataoka, H. *et al. Development Growth & Differentiation*, 39:729-740 (1997)] or the platelet endothelial cell adhesion molecule-1 (PECAM-1) (BD Pharmingen, San Diego, CA, USA, cat.no. 01951D).

[0065] PECAM-1 is a 130-kD member of the immunoglobulin (Ig) superfamily that is a major constituent of the endothelial cell intercellular junction, where up to 10^6 PECAM-1 molecules are concentrated. PECAM-1 is also expressed on the surface of circulating platelets, monocytes, neutrophils and selected T cell subsets. The tyramide signal amplification (TSA) kit (NEN Life Sciences, Boston, MA, USA) was used to enhance staining. Negative controls were produced by omitting the primary antibodies. The results were viewed with an Olympus AX80 microscope and photographed.

[0066] As can be seen from the PECAM-1 staining shown in Figures 2A-2F, the combined transfer of AdVEGF-B₁₆₇ and AdVEGF-A induced stronger formation of PECAM-1 positive vessels (Figures 2C and 2F) than AdVEGF-A alone (Figures 2B and 2E), while AdLacZ (Figures 2A and 2D) did not have any effects on the vasculature. The bar scale represents 100 micrometers.

[0067] For quantitation, the PECAM-1 positive vessels were counted at 200x magnification, using square grids (area=0.16 mm², 20x magnification). The

mean and p-values were calculated using the paired students t-test. Only values averaging higher than two standard deviations of the vessel number in the Lac virus-infected control sample were taken into consideration.

[0068] As can be seen in Figure 3, combined gene transfer of AdVEGF-B₁₆₇ and AdVEGF-A induced an approximately threefold increase in blood vessel density and VEGF-A induced a two-fold increase of blood vessel density at $p < 0.001$ in comparison to ears injected with AdVEGF-B₁₆₇ or AdLacZ alone. Thus VEGF-B potentiates the angiogenic effect of VEGF-A. It is within the scope of the invention to substitute full length, active fragments, or conservatively substituted sequences of VEGF-C or VEGF-D for VEGF-A and achieve similar results as noted above.

Example 3: Administration of Adenoviruses According to the Present Invention

[0069] A first recombinant adenovirus comprising full length or active fragments of the VEGF-B₁₆₇ sequence is prepared as described above or following other acceptable methods in the art. Conservative substitutions may be made to the sequence. A second recombinant adenovirus comprising full length or active fragments of the VEGF-A, VEGF-C, or VEGF-D sequence is prepared as described above or following other acceptable methods in the art. Conservative substitutions may be made to the sequence.

[0070] Target cells are identified for administration of the two recombinant adenovirus vectors. For example, microvascular endothelial cells may be targeted. The two vectors are administered to the target cells by methods capable of transfecting the target cells with the two vectors. For example, injection of the vectors into the microvasculature. If desired, the organism comprising the target cells may be monitored or evaluated to determine the extent of stimulation of angiogenesis.

[0071] It may be desirable to introduce the recombinant adenovirus vectors to cells proximate to the endothelial cells. Any cell type susceptible to transfection and any method of transfection may be employed. VEGFs encoded by the vectors would be secreted by the transfected cells and cause the desired angiogenic stimulation in proximate endothelial cells. For example, in the

ischemic leg model, a skeletal muscle cell could be targeted for adenovirus vector administration according to the present invention. Subsequent to transfection, the skeletal muscle cell(s) targeted would secrete the transfected VEGFs and induce cell growth and blood vessel formation in proximate endothelial cells.

[0072] The foregoing description and examples have been set forth merely to illustrate the invention and are not intended to be limiting. Since modifications of the disclosed embodiments incorporating the spirit and substance of the invention may occur to persons skilled in the art, the invention should be construed broadly to include all variations falling within the scope of the appended claims and equivalents thereof.

WHAT IS CLAIMED IS:

1. A method of stimulating angiogenesis, comprising:
introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism; and
introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-A or a fragment or conservative substitution thereof to said organism,
wherein said first replication-deficient adenovirus vector and said second replication-deficient adenovirus vector are delivered to at least one living cell selected from the group consisting of endothelial cells and cells proximate to endothelial cells of said organism.
2. A method according to claim 1, wherein said at least one living cell is a vascular cell.
3. A method according to claim 1, wherein said endothelial cells are microvascular endothelial cells
4. A method according to claim 1, wherein said endothelial cells are aortic endothelial cells.
5. A method according to one of claims 1-4, wherein said organism is mammalian.
6. A method according to one of claims 1-4, wherein said organism is murine.
7. A method according to one of claims 1-4, wherein said organism is human.

8. A method according to one of claims 1-4, wherein 10^7 to 10^{13} of vector particles of each adenovirus vector are introduced.

9. A method according to one of claims 1-4, wherein expression of said polynucleotide sequence is driven by a CMV promoter which is contained in the vector.

10. A method of stimulating angiogenesis, comprising:

—introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism; and

introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-C or a fragment or conservative substitution thereof to said organism,

wherein said first replication-deficient adenovirus vector and said second replication-deficient adenovirus vector are delivered to at least one living cell selected from the group consisting of endothelial cells and cells proximate to endothelial cells of said organism.

11. A method according to claim 10, wherein said at least one living cell is a vascular cell.

12. A method according to claim 10, wherein said endothelial cells are microvascular endothelial cells.

13. A method according to claim 10, wherein said endothelial cells are aortic endothelial cells.

14. A method according to one of claims 10-13, wherein said organism is mammalian.

15.A method according to one of claims 10-13, wherein said organism is murine.

16.A method according to one of claims 10-13, wherein said organism is human.

17.A method according to one of claims 10-13, wherein 10^7 to 10^{13} of vector particles of each adenovirus vector are introduced.

18. A method according to one of claims 10-13, wherein expression of said polynucleotide sequence is driven by a CMV promoter which is contained in the vector.

19.A method of stimulating angiogenesis, comprising:

introducing a first replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-B₁₆₇ or a fragment or conservative substitution thereof to an organism; and

introducing a second replication-deficient adenovirus vector comprising a polynucleotide sequence encoding VEGF-D or a fragment or conservative substitution thereof to said organism,

wherein said first replication-deficient adenovirus vector and said second replication-deficient adenovirus vector are delivered to at least one living cell selected from the group consisting of endothelial cells and cells proximate to endothelial cells of said organism.

20.A method according to claim 19, wherein said at least one living cell is a vascular cell.

21.A method according to claim 19, wherein said endothelial cells are microvascular endothelial cells.

22. A method according to claim 19, wherein said endothelial cells are aortic endothelial cells.

23. A method according to one of claims 19-22, wherein said organism is mammalian.

24. A method according to one of claims 19-22, wherein said organism is murine.

25. A method according to one of claims 19-22, wherein said organism is human.

26. A method according to one of claims 19-22, wherein 10^7 to 10^{13} of vector particles of each adenovirus vector are introduced.

27. A method according to one of claims 19-22, wherein expression of said polynucleotide sequence is driven by a CMV promoter which is contained in the vector.

1/3




FIG.1



FIG. 2C



FIG. 2F



FIG. 2B



FIG. 2E



FIG. 2A



FIG. 2D

3/3




FIG. 3

SEQUENCE LISTING

<110> ENHOLM, Berndt
BRANELLEC, Didier
YLAE-HERTTUALA, Seppo
PETTERSSON, Ralf
BERGSTEN, Erika
ERIKSSON, Ulf
ALITALO, Kari

<120> IN VIVO STIMULATION OF ANGIOGENIC ACTIVITY

<130> 1064/49372PC

<140> to be assigned

<141> 2001-11-01

<150> US 60/244,598

<151> 2000-11-01

<160> 2

<170> PatentIn version 3.1

<210> 1

<211> 18

<212> DNA

<213> Unknown

<220>

<223> unknown

<400> 1

cgatctggcc atacactt

18

<210> 2

<211> 25

<212> DNA

<213> Unknown

<220>

<223> unknown

<400> 2

ctatggatcc tcaccttcgc agctt

25

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/42665

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : A61K 31/70; A01N 43/04; C12N 15/63, 15/66 US CL : 514/44; 435/320.1, 455, 456 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/44; 435/320.1, 455, 456 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, DIALOG, MEDLINE, BIOSIS, EMBASE search terms: VEGF-2, VEGF-B167, angiogenesis, adenovirus, replication defective, replication deficient		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,607,918 A (ERIKSSON et al) 04 March 1997, see cols. 3-4.	1-27
Y	US 5,840,693 A (ERIKSSON et al) 24 November 1998, see cols. 3-6.	1-27
Y	US 5,792,453 A (HAMMOND et al) 11 August 1998, see col. 3, line 60 to line 26 of col. 5.	1-27
Y	US 5,652,225 A (ISNER J.M.) 29 July 1997, see cols. 1-3, and col. 6, line 13 to line 14 of col. 7.	1-27
Y	US 6,121,246 A (ISNER J.M.) 19 September 2000, see cols. 4-6.	1-27
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "G" document member of the same patent family		
Date of the actual completion of the international search 25 JANUARY 2002		Date of mailing of the international search report 13 MAR 2002
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-9230		Authorized officer QUANG NGUYEN, PH.D. Telephone No. (703) 305-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/49885

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98/32859 A1 (CORNELL RESEARCH FOUNDATION, INC.) 30 July 1998, see the entire document, especially pages 11-12 and 15-16.	1-27
Y	US 6,130,071 A (ALITALO et al) 10 October 2000, see cols. 3-17.	10-18
Y	ACHEN et al. Vascular endothelial growth factor D (VEGF-D) is a ligand for the tyrosine kinases VEGF receptor 2 (Flk1) and VEGF receptor 3 (Flt4). Proc. Natl. Acad. Sci. USA. 20 January 1998, Vol. 95, pages 548-553, see the entire document, especially page 549.	19-27
A	VERMA et al. Gene therapy---promises, problems and prospects. Nature. 18 September 1997, Vol. 389, pages 239-242.	1-27